

Research review

Self/nonself perception and recognition mechanisms in plants: a comparison of self-incompatibility and innate immunity

Author for correspondence:

Ian A. Dubery

Tel: +27 11 559 2401

Fax: +27 11 559 2401

Email: idubery@uj.ac.za

Received: 3 December 2007

Accepted: 14 January 2008

Natasha Sanabria¹, Daphne Goring², Thorsten Nürnberger³ and Ian Dubery¹

¹Department of Biochemistry, University of Johannesburg, Auckland Park, South Africa; ²Department of Cell and Systems Biology, University of Toronto, Toronto, Canada; ³Centre for Plant Molecular Biology, Eberhard-Karls University, Tübingen, Germany

Summary

Key words: innate immunity, leucine-rich repeats, pathogen-associated molecular patterns, pattern-recognition receptors, receptor-like kinases, resistance genes, S domains, self-incompatibility.

Analyses of emerging concepts indicate that parallels exist between self-incompatibility and pathogen recognition. In the case of surveillance of 'nonself', plant immune responses are triggered either by pattern recognition receptors (PRRs) that detect conserved pathogen-associated molecular patterns (PAMPs) or by resistance (R) proteins recognizing isolate-specific pathogen effectors. PAMP detection is an important component of innate immunity in plants and serves as an early warning system for the presence of potential pathogens and activation of plant defense mechanisms. In the Brassicaceae, the recognition of 'self' and self-incompatibility are components of a receptor-ligand based mechanism that utilizes an S receptor kinase (SRK) to perceive and reject 'self'-pollen. SRK is an S-domain receptor-like kinase (RLK), which in turn is part of the RLK family, some members of which represent PRRs involved in the detection of PAMPs. S-domain RLKs also occur in species that do not exhibit self-incompatibility and are up-regulated in response to wounding, PAMPs and pathogen recognition. Although evolution may have driven expansion of certain RLK families to serve roles in particular physiological processes, this may not exclude these receptor types from functioning in different programs. Recent findings on self/nonself recognition are reviewed and conceptual and mechanistic links between microbial recognition and self-incompatibility are discussed.

New Phytologist (2008) **178**: 503–514

© The Authors (2008). Journal compilation © *New Phytologist* (2008)

doi: 10.1111/j.1469-8137.2008.02403.x

Introduction

The two separate mechanisms of innate immunity and self-incompatibility (SI) are remarkably similar. The similarities

and differences between the two mechanisms in terms of functions, functional outcomes, selective processes, responses, recognition molecules, recognition receptors, and signal transduction and perception are summarized herein. In

order to elucidate how innate immunity fits into the global picture of overlapping and complex plant defense mechanisms, a short overview is presented first. In addition, an overview of SI is given to elucidate the molecular and biochemical mode of SI in the Brassicaceae. This is followed by a discussion of the role of receptor-like kinases (RLKs) in defense mechanisms and SI. While the role of S-domain RLKs in SI within the Brassicaceae is well described, the role of these receptors in pathogen perception and defense is not widely recognized.

Plant innate immunity, with its associated defense mechanisms, exhibits similar characteristics to the mammalian and insect mechanisms (Nürnberger *et al.*, 2004; Zipfel & Felix, 2005). Although they express an apparent passivity associated with their sedentary lifestyle, and are simultaneously exposed to evolving pathogens as well as environmental stresses, plants have evolved a unique metabolic plasticity that allows them to perceive pathogens and unleash effective defense strategies. The innate immune system in plants is unable to acquire or specifically adapt like the animal adaptive immune system (Goldsby *et al.*, 2000) and relies on a spectrum of predetermined receptors expressed in nonmobile cells. The question therefore arises as to how such a system could perceive so many diverse pathogen-derived signals, whilst being limited to an ancient disposition (i.e. a mechanism that originated before the evolving variables in potential invaders).

Plants therefore appear to utilize evolutionary genetic events, such as changes in gene sequence and/or genetic architecture and alterations in gene regulation, in self-defense against simultaneously evolving pathogens. Evolutionary events are also reliant on exon swapping, as well as domain recruitment through the incorporation of exons into new loci (Shapiro, 2002). Subsequently, plants must be able to re-use sections of translatable codons to produce proteins with similar morphologies or proteins that are able to multitask between different functions.

Surveillance of 'nonself': innate immunity

The ability to distinguish 'self' from 'nonself' is the most fundamental aspect of an immune system. Basal or general resistance against disease in plants used to be described by the term 'nonhost immunity', referring to an evolutionarily ancient, multilayered resistance mechanism consisting of constitutive and inducible components (Thordal-Christensen, 2003). Nonhost immunity remains operative even in susceptible plants to limit pathogen growth and is associated with the release of molecules (ligands or elicitors) derived from the pathogen and/or molecules such as oligogalacturonides and peptides released by the host plant as endogenous elicitors, analogous to the 'danger signals' of the vertebrate immune system (Matzinger, 2002). By contrast, host immunity is more recently evolved, acts within the species level and is controlled by polymorphic host genes, such as *R* (resistance)

genes, the products of which interact, directly or indirectly, with secreted 'avirulence' proteins or effectors of the pathogen (Jones & Takemoto, 2004).

The surface receptors are known to detect both pathogen-derived elicitors (or pathogen-associated molecular patterns (PAMPs) if the molecule contains a conserved 'pattern') and avirulence effectors. They include receptor-like kinases (RLKs), receptor-like proteins (RLPs) and extracellular binding proteins that may form part of multicomponent recognition complexes. Intracellular receptors are the nucleotide-binding (NB) leucine-rich repeat (LRR) class of receptors for the detection of pathogen effectors (reviewed by Nürnberger & Kemmerling, 2006; Altenbach & Robatzek, 2007; Tameling & Takken, in press, and summarized in Fig. 1).

Two branches of the plant immune system are now recognized: PAMP-triggered immunity (PTI) and effector-triggered immunity (ETI) (Chisholm *et al.*, 2006; Jones & Dangl, 2006). PTI refers to the inducible responses activated upon recognition of conserved PAMPs, such as the lipopolysaccharides (LPSs), peptidoglycan and flagellin of bacteria, and the chitin and glucan of fungi. Recent evidence indicates that some identified pattern recognition receptors (PRRs) are members of the RLK family (e.g. flagellin sensing 2 (FLS2) and the Ef-Tu receptor (EFR); Gomez-Gomez & Boller, 2000; Zipfel *et al.*, 2006). Work on flagellin and EF-Tu by the Boller group indicates that there must be a requirement for numerous such signal perception and transduction systems in plants able to recognize all potential invaders (Gomez-Gomez & Boller, 2002; Zipfel *et al.*, 2006). Sequencing of the *Arabidopsis thaliana* genome has revealed the presence of > 400 RLK sequences with various receptor configurations, of which those containing an LRR in the extracellular domain constitute the largest group, with 216 members (Shiu *et al.*, 2004; Ingle *et al.*, 2006). The diversity and large number of plant RLKs suggest that they may be involved in the perception of a wide range of stimuli (discussed in the section 'RLKs in plant innate immunity and self-incompatibility'). Other PRRs are also found amongst non-RLK proteins such as *Glycine max* beta-glucan elicitor binding proteins (GmGBP), *Lycopersicon esculentum* ethylene-inducing xylanase (LeEIX2) and chitin elicitor-binding protein (CeBIP), for perception of beta-glucans (soybean (*Glycine max*)), xylanase (tomato (*Lycopersicon esculentum*)) and chitin fragments (rice (*Oryza sativa*)), respectively (Umemoto *et al.*, 1997; Ron & Avni, 2004; Kaku *et al.*, 2006).

By contrast, ETI, the second branch, acts mostly inside the cell, using polymorphic NB-LRR proteins encoded by *R* genes. Some *R* proteins structurally resemble RLK and RLP receptors and probably evolved from PAMP receptors (reviewed by Liu *et al.*, 2007; Tameling & Takken, in press). *R* gene-mediated resistance is a form of host immunity activated upon recognition of an avirulence factor, a pathogen effector that elicits resistance, via recognition of the effector by the plant. Few *R* genes confer broad-spectrum resistance as




Fig. 1 Signaling cascades in the innate immune response of plants (adapted from Nürnberger *et al.*, 2004). Note that the non-receptor-like kinase (RLK)/receptor-like protein (RLP) pattern recognition receptors (PRRs) were not included and pathogen-associated molecular pattern (PAMP)-triggered immunity (PTI) and effector-triggered immunity (ETI) signaling pathways do not necessarily converge. (a) Pathogen recognition by the innate immune system relies on interactions between pathogen-derived molecules and corresponding host receptors. PAMPs on microbial surfaces, as well as other molecules produced by infecting pathogens, trigger innate immunity. In plants, various leucine-rich repeat (LRR)-type proteins appear to be involved in pathogen defense activation. Flagellin sensing 2 (FLS2) and the R proteins Cf9 and Xa21 are transmembrane receptors that are able to recognize PAMPs, such as flagellin, or avirulence (Avr) effector signals. Avr9 is recognized by a high-affinity binding site in tomato. This complex interacts with Cf9 and activates at least two mitogen activated protein kinase (MAPKs). *Arabidopsis thaliana* FLS2 and rice Xa21 are likely to transduce the pathogen signal through their cytoplasmic protein kinase domains. The amino-terminal fragment of flagellin (flg22) directly binds to FLS2 and activates MAPKs, AtMPK3 and AtMPK6. Translocation of PAMP-activated plant MAPKs into the nucleus has been demonstrated, where these enzymes are likely to contribute to the activation of transcription factors of the WRKY type. In turn, intracellular plant R proteins recognizing Avr signals confer pathogen race/plant cultivar-specific immunity to viral (tobacco resistance gene to tobacco mosaic virus (N) and potato resistance gene to potato virus X (Rx)), bacterial (*Arabidopsis* resistance gene to *Pseudomonas syringae* pv tomato strain DC3000 (RPS4), *Arabidopsis* resistance gene to *Pseudomonas syringae* pv tomato strains producing AvrRpm1 or AvrB (RPM1) and *Arabidopsis* resistance gene to *Pseudomonas syringae* pv tomato strains that produce AvrRpt2 (RPS2)), oomycete (*Arabidopsis* resistance gene to *Peronospora parasitica* (RPP5)), or fungal (flax rust resistance gene (L⁶)) pathogens (Gomez-Gomez & Boller, 2002). Intracellular nucleotide binding site (NBS)-LRR proteins are linked to coiled-coil (CC) or Toll-interleukin 1 receptor-like (TIR) domains. PGN, peptidoglycans; Pto, a tomato protein kinase. (b) Ligand-induced dimerization. During the two-step address-message mechanism, binding of flagellin to the N-terminal part is the first step and activation of responses with the C-terminus is the second step (Meindl *et al.*, 2000). flg22-dependent heterodimerization of FLS2 and brassinosteroid receptor-associated receptor kinase (BAK1) occurs, where FLS2 binds flg22 independently of its association with BAK1. BAK1 probably does not determine the specificity of the signal output, but is likely to have a role as an adaptor or coreceptor for regulation of various receptors, as suggested by Chinchilla *et al.* (2007).

they act in a race-specific manner. *R* gene-mediated immunity is often associated with the hypersensitive response (HR). It results in local induced resistance (LIR), acting at the site of infection to contain the invader, and systemic acquired

resistance (SAR), which induces defenses in distal, noninfected parts of plants after activation of local resistance. It should be noted that SAR has also been demonstrated to be induced by PAMP recognition (Mishina & Zeier, 2007).




Fig. 2 S receptor kinase (SRK) signaling in the *Brassica* self-incompatibility response (adapted from Haffani *et al.*, 2004; Takayama & Isogai, 2005). (a) Inactive, but primed. Ligand-independent dimerization might provide a 'primed' state that allows rapid recruitment and activation of the receptor on ligand binding (Naithani *et al.*, 2007). In the absence of incompatible pollen, SRK is inhibited by thioesteronin *h* proteins (THL1/2). This inhibition is released upon the haplotype-specific S cysteine-rich (SCR)/S locus protein 11 (SP11) pollen ligand binding (Bower *et al.*, 1996; Cabrilla *et al.*, 2001). S locus glycoprotein (SLG) and a soluble extracellular domain produced from SRK (eSRK) represent low-affinity binding sites for SCR/SP11. (b) Active. During incompatible pollination, the SCR/SP11 ligand binds to and activates SRK (Giranton *et al.*, 2000). Whether SLG participates in the SRK complex is unclear, as the haplotype-specific SLG is not always required. Activated SRK is proposed to autophosphorylate serine and threonines, and some of these phosphorylation sites represent docking sites for downstream signaling proteins such as armadillo-repeat-containing 1 (ARC1) and the M-locus protein kinase (MLPK).

Dangl and Jones have questioned the likelihood of the known *R* genes, which are linked to defense, being able to recognize all the possible effector signals (Dangl & Jones, 2001). The mechanisms of *R* gene-mediated immunity may thus be explained by the 'gene-for-gene' genetic model or the 'guard hypothesis' molecular model. A 'guard' can refer to a typical *R* protein, whereas the 'guardee' represents a target of pathogen effectors (Dangl & Jones, 2001). Many plant *R* proteins might be activated indirectly by pathogen-encoded effectors, and not by direct recognition (Dangl & Jones, 2001). This form of 'guard hypothesis' implies that *R* proteins are able to indirectly recognize pathogen effectors by monitoring the structural integrity of the host cell targets, which is altered by effector action. The *R* proteins in question are thus activated as sensors of 'pathogen-induced self' or 'altered self' molecular patterns and can potentially perceive the presence of more than one effector protein. This can explain how plants can potentially recognize a diverse set of pathogens and pathogen-specific molecules, using a relatively limited number of pathogen receptors.

A 'zigzag' model to illustrate the quantitative output of the plant immune system, as well as to illustrate the evolutionary relationship between PTI and ETI, was recently proposed (Jones & Dangl, 2006). In phase 1, PAMPs or microbe-associated molecular patterns (MAMPs) (i.e. pathogens or microbes) are recognized by PRRs, resulting in PTI, which can stop further

colonization. In phase 2, successful pathogens deploy effectors that contribute to pathogen virulence. When effectors interfere with PTI, this results in effector-triggered susceptibility (ETS). In phase 3, an effector is specifically recognized by an *R* protein, which results in ETI. Therefore, recognition is either indirect, or through direct NB-LRR recognition of an effector. ETI is an accelerated and amplified PTI response, which results in disease resistance and may lead to the HR at the infection site. In phase 4, natural selection drives pathogens to avoid ETI. This is achieved either by shedding or diversifying the recognized effector gene, or by acquiring effectors that suppress ETI. Thereafter, natural selection results in the evolution of new *R* specificities leading to ETI being triggered again.

Transcriptional analysis of genes expressed in *A. thaliana* in response to elicitation by flagellin (flg22) (Navarro *et al.*, 2004) indicates that a considerable number of the up-regulated genes can be classified as being involved in signal perception (*RLK* and *R* genes) and signal transduction. This is indicative of positive feedback regulation operating in innate immunity with transcriptional activation of the components involved in the perception and signaling (Zipfel *et al.*, 2004). The up-regulated expression of *RLK* and *R* genes presumably leads to an enhanced sensitivity of the plant to further stimuli, which allows sensing of the presence of invading microorganisms with other PAMPs or effector signals; that is, a primed or sensitized state.

Surveillance of 'self': self-incompatibility

The established system of 'self'/'nonself' recognition in SI systems utilizes receptor–ligand type interactions to perceive, recognize and reject incompatible pollen. Thus, SI prevents 'self'-pollination (Fig. 2). Although SI responses are generally comprised of a 'self' and 'nonself' recognition process, SI systems have evolved independently and do not utilize one molecular mechanism exclusively. Rather, SI encompasses a collection of divergent cellular responses leading to pollen rejection (Takayama & Isogai, 2005; Wheeler & Franklin-Tong, 2007).

The molecular signatures of 'self' and 'nonself' in plant SI are unambiguous. The recognition and rejection of 'self'-pollen is generally regulated by two or more multiallelic and tightly linked *S* genes (comprising *S* haplotypes). In sporophytic systems such as in the Brassicaceae, 'self' is derived from the expression of matched products from the same *S* haplotype in the interacting pistil and pollen parent, whereas 'nonself' is derived from the expression of the unmatched products from different *S* haplotypes (Takayama & Isogai, 2005). This system uses a receptor–ligand based mechanism, with the S-domain receptor kinase (SRK), to perceive a ligand, the S cysteine-rich (SCR)/*S* locus protein 11 (SP11), present on the pollen coat. The multiallelic *SRK* gene is the 'female' determinant of specificity in the SI response of the Brassicaceae (Takasaki *et al.*, 2000; Silva *et al.*, 2001). SRK spans the plasma membrane of stigmatic epidermal cells, and it is activated in an *S* haplotype-specific manner upon binding of the pollen ligand to a hypervariable subdomain in its extracellular region (Fig. 2; Kachroo *et al.*, 2001; Takayama *et al.*, 2001; Kemp & Doughty, 2007; Shimosato *et al.*, 2007). The multiallelic *SCR/SP11* gene is the 'male' determinant in this system (Schopfer *et al.*, 1999; Takayama *et al.*, 2000). Activation of SRK leads to cellular signaling pathways in the stigmatic papillae causing a block in pollen hydration, germination and pollen tube growth.

Another protein implicated in the *Brassica* SI system is the multiallelic *S* locus glycoprotein (SLG), a secreted glycoprotein encoded in the *S* locus region and expressed in the stigma. SLG shows a sequence similarity to the ectodomain of SRK, but does not contribute to the *S* haplotype specificity of this system (Takasaki *et al.*, 2000; Silva *et al.*, 2001). The expression of SLG has been proposed to facilitate the processing or accumulation of SRK (Dixit *et al.*, 2000), or to enhance the SI response (Takasaki *et al.*, 2000), although it is not always required (Silva *et al.*, 2001). SLG was co-immunoprecipitated as part of a chemically cross-linked SRK complex and may function as a coreceptor with SRK to form a heteromeric receptor complex that perceives the signal carried by the pollen (Giranton *et al.*, 2000). However, SLG and a soluble extracellular domain produced from SRK (eSRK) primarily exist as monomers. Recently, the soluble SLG and eSRK were found only to present low-affinity binding sites for SCR/SP11, while the membrane-bound SRK and a truncated

membrane-bound form of SRK (tSRK) presented high-affinity binding sites for SCR/SP11 (Shimosato *et al.*, 2007). While tSRK could participate in complexes with SRK for high-affinity SCR/SP11 binding, it is less likely that SLG and eSRK participate in these complexes.

The classical view of ligand-activated animal receptors involves receptor homodimerization or oligomerization, which is induced by ligand binding and serves to bring the receptor intracellular domains into close proximity for transphosphorylation and recruitment of effector cytoplasmic proteins (Heldin, 1995). SRK has been found to exist as a dimer in unpollinated pistils (i.e. in the absence of ligand), suggesting that the classical animal-based receptor model does not apply (Fig. 2; Giranton *et al.*, 2000). In this state, the SRK dimer provided a high-affinity binding site for the *S* haplotype-specific SCR/SP11 ligand, and, interestingly, high-affinity binding of SCR/SP11 appears to be a consequence of the presence of preformed dimers (Shimosato *et al.*, 2007). The ligand-independent dimerization of SRK might provide a 'primed' condition that allows the rapid recruitment and activation of the receptor on ligand binding. SCR/SP11 ligand binding may then cause the rearrangement of existing SRK dimers or stabilize the ligand–receptor complex, leading to the phosphorylation and activation of SRK (Shimosato *et al.*, 2007). The recruitment of this additional step for full receptor activation has also been reported for animal receptors (Heldin, 1995; Giranton *et al.*, 2000).

Self-incompatibility vs innate immunity

Plant SI and plant immunity evolved in response to different pressures, namely, avoidance of inbreeding in the former case and avoidance of parasitism in the latter. The recognition and rejection of 'self' in plant SI can be compared to recognition and defense activation in plants. The *Brassica* SI SRK complex serves to recognize, and mounts a response to, 'self' ligands. By contrast, in plant defense, pathogen-derived 'nonself' ligands are recognized by RLKs and pathogen-induced changes to 'guard' molecules are recognized as 'altered self' by *R* gene products.

Hogenboom (1983) noted the close parallels between the genetics of SI and plant–pathogen interactions. Hodgkin *et al.* (1988) compared SI responses with pathogen recognition and pointed out that parallels between SI and host–pathogen interactions include the penetration of the 'host' by a tubular cell emanating from a spore-like structure. Support for this idea came later when the wheat (*Triticum aestivum*) wheat leaf rust kinase (*WLRK*) defense genes were found to be structurally related to *SRK* genes, and this led to speculation that genes involved in SI and defense might have had a common ancestor (Feuillet *et al.*, 1998). Nasrallah (2005) has discussed the evolutionary origins of plant SI, focusing on the hypothesis that SI evolved from a defense pathway (Hiscock *et al.*, 1996).

Table 1 Summarized comparison between plant innate immunity and self-incompatibility

	Innate immunity	Self-incompatibility (specific reference to Brassicaceae)
Physiological function	Basal defense	Reproduction
Selective process	Recognition of: nonself: (PAMP), PTI altered self: (Avr-R), ETI, endogenous elicitors	Recognition of: self
Response	Mounts response to nonself or altered self	Mounts response to self
Recognition molecule	PAMPs Avr effectors	SCR protein ligand
Recognition receptor	Host-derived oligouronides and peptides RLK PRRs (e.g. for PAMPs) R proteins (directly or indirectly, e.g. for pathogen effectors) non-RLK PRRs (e.g. for endogenous oligogalacturonides, glucans, or chitin fragments)	S-domain RLKs for pollen epitopes
Receptor activation	PRRs bind to PAMPs with or without coreceptor	S-domain RLKs form oligomers and bind to pollen
Subsequent reactions during response	Downstream intracellular phosphorylation cascades are triggered Deposition of callose at site of plant cell in host–pathogen interaction Cross protection; an incompatible host–pathogen interaction can affect the outcome of a compatible interaction Induction of general inhibitory compounds, e.g. phytoalexins Ca ²⁺ -dependent signaling network Programmed cell death	Downstream intracellular signaling cascades are triggered Deposition of callose at surface of stigma cells Individual interaction; an individual pollen grain interacts with an individual papillar cell No induction of general inhibition that would interfere with compatible pollen reactions Ca ²⁺ -dependent signaling network ¹ DNA fragmentation and morphological changes in mitochondria, Golgi and endoplasmic reticulum ¹
Functional outcome	Plant rejects pathogen by block in pathogen penetration and proliferation	Plant rejects pollen by block in pollen hydration, tube penetration and growth

¹As seen in the poppy system, but not in *Brassica*.

Avr, avirulence; Avr-R, avirulence-resistance; ETI, effector-triggered immunity; PAMP, pathogen-associated molecular pattern; PRR, pattern recognition receptor; PTI, PAMP-triggered immunity; RLK, receptor-like kinase; SCR, S cysteine-rich.

Parallels exist where plant SI and plant immunity have similar outcomes, such as the elimination of undesirable cells or organisms. Also, both immunity and at least some SI systems (the crucifer and possibly the poppy systems) use highly variable receptors to recognize highly variable ligands. In addition, SI systems that have bio-destructive activity towards pollen tubes use components that are also used in defense; for example, programmed cell death and other reactions triggered in the incompatible pollen tubes of poppy are also induced during the plant immune response (Jordan *et al.*, 2000; Dangl & Jones, 2001; Geitmann *et al.*, 2004; Thomas & Franklin-Tong, 2004).

Another example relates to the nature of SCR/SP11, the pollen determinant of SI specificity in crucifers. SCR/SP11 is similar in structure, although not in primary sequence, to

defensins, the molecules of innate immunity that present a first line of defense to microbial challenge in plants and animals (Mishima *et al.*, 2003; Chookajorn *et al.*, 2004). The similarity between the two classes of molecules suggests an evolutionary link, albeit a distant one, between crucifer SI and innate immunity.

The most notable parallels, however, emerge from comparisons of the self-recognition loci and genes of plant SI with those that control self/nonself recognition in a variety of recognition systems, the vertebrate major histocompatibility complex (MHC) in particular (Janeway & Medzhitov, 2002). In both plants and animals, and from immunity to reproduction, self/nonself discrimination systems have been molded by similar selective pressures for diversification and coevolution of recognition functions, and by a shared requirement to maintain the genetic linkage of coadapted gene complexes (Nasrallah, 2005).

Table 2 Summary of putative associations of S-domain receptor-like kinases (RLKs) with defense mechanisms

Gene	Accession number	Organism	Association	Reference
ARK1	gi/18408364	<i>Arabidopsis thaliana</i>	Bacteria and wounding inducible	Pastuglia <i>et al.</i> (2002)
ARK3	gi/30685418	<i>Arabidopsis thaliana</i>	Bacteria and wounding inducible	Pastuglia <i>et al.</i> (2002)
HAP3–15	gi/67568666/gb/DR109311.1	<i>Nicotiana tabacum</i>	Putative S-domain receptor-like kinase with protein–protein or protein–carbohydrate interactions	Sanabria & Dubery (2006)
<i>Pi-d2</i>		<i>Oryza sativa</i>	<i>R/RLK</i> gene with extracellular S domain	Chen <i>et al.</i> (2006)
RKS1	gi/4008007/gb/AF084035.1	<i>Arabidopsis thaliana</i>	Salicylic acid inducible	Takahashi <i>et al.</i> (1998)
RKS2	gi/4008009/gb/AF084036.1	<i>Arabidopsis thaliana</i>	Salicylic acid inducible	Takahashi <i>et al.</i> (1998)
RLK1	gi/18424408/NM_125483	<i>Arabidopsis thaliana</i>	Salicylic acid inducible	Walker (1993)
SFR1	gi/2598268/emb/Y14285.1	<i>Brassica oleracea</i>	Bacteria and salicylic acid inducible	Pastuglia <i>et al.</i> (2002)
SFR2	gi/1783311/emb/X98520.1	<i>Brassica oleracea</i>	Bacteria and salicylic acid inducible	Pastuglia <i>et al.</i> (2002)
SI-RLK1	gi/146739162/EF560751	<i>Oryza sativa</i>	Salt/stress inducible	Unpublished, direct submission to NCBI
	At5g60900	<i>Arabidopsis thaliana</i>	S-receptor kinase homolog 2 precursor	T. Nürnberger (unpublished)*
	At5g18470	<i>Arabidopsis thaliana</i>	Putative protein S-receptor kinase PK3 precursor	T. Nürnberger (unpublished)*
	At1g70530	<i>Arabidopsis thaliana</i>	Putative protein kinase similar to C-terminal region of S-receptor kinase precursor	T. Nürnberger (unpublished)*

*TAIR accession expression set 100808727. *ARK1*, *Arabidopsis* receptor kinase 1; *ARK3*, *Arabidopsis* receptor kinase 3; *HAP3–15*, *Hind* III arbitrary primer; *Pi-d2*, resistance gene *Pi-d(t)2*, renamed as *Pi-d2*, confers resistance to the *M. grisea* strain ZB15; *RKS1*, receptor-like protein kinase 1; *RKS2*, receptor-like protein kinase 2; *RLK1*, receptor-like protein kinase; *SFR1*, S family receptor 1; *SFR2*, S family receptor 2; *SI-RLK1*, stress-induced receptor-like kinase 1.

In this context, the similarities between plant SI and plant innate immunity have, however, received scant attention. Table 1 summarizes the major similarities and differences between innate immunity and SI in plants with regard to functions, functional outcomes, selective processes, responses, recognition molecules, recognition receptors, and signal transduction and perception.

RLKs in plant innate immunity and self-incompatibility

The proposed evolutionary relationships among receptor kinase family members arose from an ancient duplication event leading to the divergence of RLK/Pelle from the receptor tyrosine kinase (RTK)/Raf group, which consists of serine/threonine kinases. Thereafter, a more recent gene duplication led to the divergence of RTK from Raf, followed by the divergence of plant and animal lineages, resulting in the ancestral sequences that gave rise to the extant receptors and related kinases (Shiu & Bleeker, 2001). The evolutionary history of plant RLKs indicates that the kinase domains were recruited numerous times by fusion with different extracellular domains to form the subfamilies found in *A. thaliana*. Based on the presence or absence of extracellular domains, members of this gene family are categorized as RLKs or receptor-like cytoplasmic kinases (RLCKs). Subfamilies are assigned based on kinase phylogeny and are grouped according to the domain organization of the majority of members in a given subfamily (comparative summary reported in Shiu & Bleeker, 2001).

It has been suggested that a drastic expansion of the *RLK* gene family occurred in the land plant lineage and that this abundance of plant RLKs represents a plant-specific utilization for extracellular signal sensing. Diverse sequence motifs are present in the extracellular domains of RLKs (Shiu & Bleeker, 2001) and these motifs are potentially responsible for interactions with other proteins, carbohydrates or lipids. The data indicate that RLKs involved in resistance or defense responses may have been duplicated or retained at higher rates in a lineage-specific fashion (Shiu *et al.*, 2004). The preferential expansion of defense/resistance-related RLKs could be the consequence of strong selection pressure for recognizing pathogens (Shiu *et al.*, 2004). The large family of plant RLK proteins therefore contains distinct protein kinases, each of which might play a unique role in cellular signaling (Walker, 1994; Haffani *et al.*, 2004), and probably comprise receptors for further PAMP recognition (Zipfel *et al.*, 2006).

An example of independent recruitment of biochemical components for different functions is the LRR motif. LRR domains are found in transmembrane proteins, transmembrane kinases and intracellular R proteins. Collectively, LRRs appear to be involved in a range of processes from development to intercellular communication and disease resistance (Zhang, 1998; Torii, 2004; Chisholm *et al.*, 2006). A number of LRR transmembrane and intracellular proteins act as integral components of ligand perception complexes during ETI (Dangl & Jones, 2001). In addition, the LRR motif also plays an important role in PRRs in the evolutionarily older PTI (Nürnberger & Kemmerling, 2006). Although leucine-rich repeat RLKs (LRR-RLKs) (particularly the members of

'clade XII') have been implicated in plant immunity, and the S-domain RLKs (particularly SRK) have been associated with SI, this does not exclude the possibility that other receptor types are involved in either program. The role of LRR-RLKs in defense mechanisms is now recognized (e.g. FLS2, EFR and the endogenous peptide ligand of the AtPEP 1 receptor (PEPR1): Gomez-Gomez & Boller, 2000; Yamaguchi *et al.*, 2006; Zipfel *et al.*, 2006), but intermittent reports on the potential involvement of S-domain RLKs in defense mechanisms (Table 2) have received little attention (Pastuglia *et al.*, 1997, 2002; Bassett *et al.*, 2005).

S-domain *RLK* genes belong to large subfamilies with 40 and 147 members in *A. thaliana* and rice, respectively (Morillo & Tax, 2006). *RLK* genes containing the S domain are regarded as unique intermediates between RLKs mediating developmental and resistance functions (Shiu & Bleecker, 2001; Shiu *et al.*, 2004). Except in the case of the SRK involved in *Brassica* SI, very little is known about their functions. It has been suggested that the development of the SRK-mediated SI response is an evolutionarily relatively recent event in the Brassicaceae and may have occurred through the recruitment of pre-existing genes that performed other related functions (Shiu & Bleecker, 2001). The widespread occurrence of S-domain RLKs in *A. thaliana* and other plants, expressed in nonreproductive tissues, and in species that do not exhibit SI (Pastuglia *et al.*, 2002; Bassett *et al.*, 2005), would also argue for additional functions for S-domain RLKs. This leads to the questions of whether the conserved S domain has a function other than in SI and whether the SI-linked SRKs have an earlier S-domain ancestor, linked to defense.

S-domain RLKs, like other RLKs, contain all the elements required for PAMP perception and signal transduction: the proteins are single-pass transmembrane serine/threonine kinases displayed on the plasma membrane. The extracellular domains of S-domain RLKs include a mannose-binding agglutinin/B-lectin domain, a cysteine-rich S domain and an epidermal growth factor (EGF)-like or plasminogen/apple/nematode (PAN) motif (Tordai *et al.*, 1999; Shiu & Bleecker, 2001). The PAN motif represents a conserved module, also found in the ectodomain of several animal receptors, that functions in protein–protein interactions, similarly to LRR domains, but also in protein–carbohydrate (e.g. mannose binding) interactions (Tordai *et al.*, 1999). The region containing the predicted PAN motif plays a primary role in dimerization of the receptor, and the S-domain motif plays a secondary role. The extracellular domain of SRK showed a preference, mediated by a small, highly variable region within the PAN motif, for homodimers over heterodimers with the products of other SRK alleles. Thus, the polymorphic extracellular domain of SRK is not only responsible for *S* haplotype-specific binding of the SCR/SP11 ligand (Kemp & Doughty, 2007), but also appears to play a role in the allele-specific homodimerization of SRK (Naithani *et al.*, 2007).

Another feature possibly exhibited by S-domain RLKs (Giranton *et al.*, 2000), and LRR-RLKs involved in PAMP perception, is the involvement of a coreceptor that may modulate specificity (Chinchilla *et al.*, 2007; Heese *et al.*, 2007; see legend to Fig. 1b). As discussed in the section 'Surveillance of "self": self-incompatibility', oligomeric complexes of receptors with bound ligands lead to transphosphorylation of the receptor and triggering of a signaling cascade (Figs 1, 2). Signal transduction downstream of RLKs involves proteins that interact with and are phosphorylated by the cytoplasmic domain of the RLK. In the case of the *Brassica* SI system, the plant-specific signaling protein armadillo-repeat-containing 1 (ARC1) binds to SRK (Gu *et al.*, 1998; Stone *et al.*, 1999). The ARC1 protein belongs to the U-box family of E3 ligases, and, interestingly, other related members have been implicated in plant cell death and defence responses in *A. thaliana*, tobacco (*Nicotiana tabacum*) and rice (Zeng *et al.*, 2004; Gonzalez-Lamothe *et al.*, 2006; Yang *et al.*, 2006). A reduced expression of *ARC1* led to a partial breakdown of SI, providing evidence that ARC1 acts immediately downstream of SRK as a positive modulator of SI (Stone *et al.*, 1999). A U-box in ARC1 suggests a role for ubiquitination in the SI response (Newbiggin & Vierstra, 2003). Ubiquitin is well known in defense mechanisms as the tag that directs targeted proteins to the machinery that eliminates them (Devoto *et al.*, 2003). Phosphorylated ARC1 relocates from the cytosol to the proteasome present on the cytosolic face of endoplasmic reticulum membranes and promotes the ubiquitination and proteasomal degradation of unknown substrate proteins, thereby leading to pollen rejection (Stone *et al.*, 2003).

SRK may also interact with other signal transduction-associated molecules such as calmodulin and the kinase-associated protein phosphatase (KAPP; Vanoosthyse *et al.*, 2003). It should be noted that FLS2 also interacts with KAPP (Gómez-Gómez *et al.*, 2001; Ding *et al.*, 2007). In addition, SRK interacts with the M-locus protein kinase, a member of the RLCK family, that is required downstream of SRK to promote the SI response and is efficiently phosphorylated by SRK *in vitro* (Murase *et al.*, 2004; Kakita *et al.*, 2007).

A number of S-domain *RLK* genes have been found to have up-regulated expression in response to pathogen infections, wounding, or treatment with defence-related compounds such as salicylic acid (a metabolite that plays an important role in potentiating local and systemic induced resistance) and LPS (Table 2). Interestingly, the first report of salicylic acid-induced expression of *RLK* genes was found in a member of the *Brassica* S-domain RLK family, *S* family receptor 2 (*SFR2*). *SFR2* expression was found to be induced by bacterial pathogens, wounding, and treatment with salicylic acid. The transient induction of the *SFR2* gene exhibited a kinetic and induction pattern typical of defense genes (Pastuglia *et al.*, 1997). A second closely related S-domain *RLK* gene, *SFR1*, was also found to be induced by bacterial infection and salicylic acid treatment (but not wounding) in *Brassica oleracea* (Pastuglia *et al.*, 2002).

In addition, two closely related *A. thaliana* S-domain *RLK* genes, *Arabidopsis* receptor kinase (*ARK*)1 and *ARK*3, were found to have increased mRNA accumulation following bacterial infections and wounding (Pastuglia *et al.*, 2002). Two other *A. thaliana* S-domain *RLK* genes, receptor-like protein kinase (*RKS*)1 and *RKS*2, are also induced by salicylic acid treatment (Ohtake *et al.*, 2000). Finally, it has been reported that LPS elicitation of *N. tabacum* cell suspensions resulted in the differential expression of a putative S-domain *RLK* (Sanabria & Dubery, 2006). In addition, three S-domain *RLK* genes (At5g60900, At5g18470 and At1g70530) were found to be up-regulated in a transcriptional microarray analysis of genes expressed in *A. thaliana* in response to elicitation by LPS (T. Nürnberger, unpublished: TAIR accession expression set 100808727), suggesting a putative function in LPS perception.

What is probably the first direct genetic evidence for the role of S-domain receptor kinases in plant disease resistance comes from Chen *et al.* (2006), who characterized an *R* gene (Pi-d2) that confers resistance to blast disease in rice. The gene encodes a bulb-type mannose-specific binding (B)-lectin receptor kinase and belongs to the S-domain-2b *RLK* subfamily of lectin receptor kinases (LecRKs). Although several *R* genes that encode *RLK*s have been cloned and characterized, this is the first reported to have an extracellular lectin domain and to belong to the S-domain *RLK*s. Despite the presence of this domain, an indirect role for Pi-d2 in pathogen recognition was proposed without considering the variable features of the extracellular domain of S-domain *RLK*s that allow them to multitask in developmental and defense responses.

A summary of S-domain *RLK*s and their association with defense mechanisms is given in Table 2.

Conclusions

In addition to PAMP perception as a 'nonself' surveillance mechanism in innate immunity, the evolutionary solution in plants to the problems of perceiving and responding to pathogens involves 'self' surveillance, which is conceptually similar to SI. The two separate mechanisms of innate immunity and SI are remarkably similar, leading to speculation about a common ancestor for genes involved in SI and defense and the hypothesis that SI evolved from a defense pathway.

Similar to the animal innate immune system (Medzhitov & Janeway, 2002), different and possibly overlapping receptor types may be implicated in different physiological programs in plants, such as immunity and SI. There is currently no conclusive evidence for evolutionary conservation of an ancient PAMP detection system (Dangl & Jones, 2001) and independent recruitment of components during evolution is equally plausible (Ingle *et al.*, 2006). Moreover, there are also various examples where a specific type of biochemical module or protein appears to be used to fulfil a requirement in more than one process, that is, to show dual functioning,

and the re-use of highly evolved processes for diverse functions was recently pointed out by Ausubel (2005) in his perspective on immune signaling pathways. There are thus some indications that certain defense genes are structurally related to the S-domain *RLK* genes which can be regarded as intermediates between *RLK* genes mediating developmental and resistance functions. Specific domains in *RLK* proteins can be utilized to fulfil a number of biochemical functions and receptor modules are not necessarily reserved for one physiological purpose only. Although evolution may have driven expansion of particular *RLK* families (LRR-*RLK*s and S-domain *RLK*s) to serve roles in particular physiological processes (defense/development and SI, respectively), this may not exclude these receptor types from functioning in different programs, lending support to the hypothesis that subsets of molecules involved in innate immunity were co-opted to perform 'self' recognition functions in reproduction. It is thus plausible that S-domain *RLK* genes could be utilized to function as *R* genes or as PRRs in perception of PAMPs of a nonprotein nature.

Unfortunately, the role of S-domain *RLK*s in defense mechanisms has previously not been widely recognized, or thoroughly explored. Further research is therefore warranted in order to broaden our understanding of the involvement or dual-functioning of S-domain *RLK*s in PAMP surveillance and perception. This includes aspects of receptor function and ligand–receptor interaction, the sharing of receptors between ligands with common molecular signatures and the modulating role of potential co-receptors in interaction, specificity and priming.

References

- Altenbach D, Robatzek S. 2007. Pattern recognition receptors: from the cell surface to intracellular dynamics. *Molecular Plant–Microbe Interactions* 20: 1031–1039.
- Ausubel FM. 2005. Are innate immune signalling pathways in plants and animals conserved? *Nature Immunology* 6: 973–979.
- Bassett CL, Nickerson ML, Farrel RE, Artlip TS, Ghaouth AE, Wilson CL, Wisniewski ME. 2005. Characterization of an S-locus receptor protein kinase-like gene from peach. *Tree Physiology* 25: 403–411.
- Bower MS, Matias DD, Fernandes-Carvalho E, Mazzurco M, Gu T, Rothstein SJ, Goring DR. 1996. Two members of the thioredoxin-h family interact with the kinase domain of a *Brassica* S locus receptor kinase. *Plant Cell* 8: 1641–1650.
- Cabrillac D, Cock JM, Dumas C, Gaude T. 2001. The S-locus receptor kinase is inhibited by thioredoxins and activated by pollen coat proteins. *Nature* 410: 220–223.
- Chen X, Shang J, Chen D, Lei C, Zou Y, Zhai W, Liu G, Xu J, Ling Z, Cao G *et al.* 2006. A B-lectin receptor kinase gene conferring rice blast resistance. *Plant Journal* 46: 794–804.
- Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nürnberger T, Jones JDG, Felix G, Boller T. 2007. A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defense. *Nature* 448: 497–501.
- Chisholm ST, Coaker G, Day B, Staskawicz BJ. 2006. Host–microbe interactions: shaping the evolution of the plant immune response. *Cell* 124: 803–814.
- Chookajorn T, Kachroo A, Ripoll DR, Clark AG, Nasrallah JB. 2004. Specificity determinants and diversification of the *Brassica*

- self-incompatibility pollen ligand. *Proceedings of the National Academy of Sciences, USA* 101: 911–917.
- Dangl JL, Jones JDG. 2001. Plant pathogens and integrated defense responses to infection. *Nature* 411: 826–833.
- Devoto A, Muskett PR, Shirasu K. 2003. Role of ubiquitination in the regulation of plant defence against pathogens. *Current Opinions in Plant Biology* 6: 307–311.
- Ding Z, Wang H, Liang X, Morris ER, Gallazzi F, Pandit S, Skolnick J, Walker JC, Van Doren SR. 2007. Phosphoprotein and phosphopeptide interactions with the FHA domain from Arabidopsis kinase-associated protein phosphatase. *Biochemistry* 46: 2684–2696.
- Dixit R, Nasrallah ME, Nasrallah JB. 2000. Post-transcriptional maturation of the S receptor kinase of *Brassica* correlates with co-expression of the S-locus glycoprotein in the stigmas of two *Brassica* strains and in transgenic tobacco plants. *Plant Physiology* 124: 297–311.
- Feuillet C, Reuzeau C, Kjellbom P, Keller B. 1998. Molecular characterization of a new type of receptor-like kinase (WLRK) gene family in wheat. *Plant Molecular Biology* 37: 943–953.
- Geitmann A, Franklin-Tong VE, Emons AC. 2004. The self-incompatibility response in *Papaver rhoeas* pollen causes early and striking alterations to organelles. *Cell Death and Differentiation* 11: 812–822.
- Giranton J-L, Dumas C, Cock JM, Gaude T. 2000. The integral membrane S-locus receptor kinase of *Brassica* has serine/threonine kinase activity in a membranous environment and spontaneously forms oligomers in planta. *Proceedings of the National Academy of Sciences, USA* 97: 3759–3764.
- Goldsby RA, Kindt TJ, Osborne BA. 2000. *Kuby immunology*, 4th edn. New York, NY, USA: W. H. Freeman & Co.
- Gómez-Gómez L, Bauer Z, Boller T. 2001. Both the extracellular leucine rich repeat domain and the kinase activity of FLS2 are required for flagellin binding and signalling in Arabidopsis. *Plant Cell* 13: 1155–1163.
- Gomez-Gomez L, Boller T. 2000. FLS2, an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in Arabidopsis. *Molecular Cell* 5: 1003–1011.
- Gomez-Gomez L, Boller T. 2002. Flagellin perception: a paradigm for innate immunity. *Trends in Plant Science* 7: 251–256.
- Gonzalez-Lamothe R, Tsitsigiannis DI, Ludwig AA, Panicot M, Shirasu K, Jones JD. 2006. The U-box protein CMPG1 is required for efficient activation of defense mechanisms triggered by multiple resistance genes in tobacco and tomato. *Plant Cell* 18: 1067–1083.
- Gu T, Mazzurco M, Sulaman W, Maltias DD, Goring DR. 1998. Binding of an arm repeat protein to the kinase domain of the S-locus receptor kinase. *Proceedings of the National Academy of Sciences, USA* 95: 382–387.
- Haffani YZ, Silva NF, Goring DR. 2004. Receptor kinase signalling in plants. *Canadian Journal of Botany* 82: 1–15.
- Heese A, Hann DR, Gimenez-Ibanez S, Jones AME, He K, Li J, Schroeder JI, Peck SC, Rathjen JP. 2007. The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants. *Proceedings of the National Academy of Sciences, USA* 104: 12217–12222.
- Heldin CH. 1995. Dimerization of cell surface receptors in signal transduction. *Cell* 80: 213–223.
- Hiscock SJ, Kües U, Dickinson HG. 1996. Molecular mechanisms of self-incompatibility in flowering plants and fungi – different means to the same end. *Trends in Cell Biology* 6: 421–428.
- Hodgkin T, Lyon DG, Dickinson HG. 1988. Recognition in flowering plants: a comparison of the *Brassica* self-incompatibility system and plant pathogen interactions. *New Phytologist* 110: 557–569.
- Hogenboom NG. 1983. Bridging the gap between related fields of research: pistil-pollen relationships and the distinction between incompatibility and incongruity in nonfunctioning host-parasite relationships. *Phytopathology* 73: 381–383.
- Ingle RA, Carstens M, Denby KJ. 2006. PAMP recognition and the plant-pathogen arms race. *Bioessays* 28: 880–889.
- Janeway CA, Medzhitov R. 2002. Innate immune recognition. *Annual Review of Immunology* 20: 197–216.
- Jones DA, Takemoto D. 2004. Plant innate immunity – direct and indirect recognition of general and specific pathogen-associated molecules. *Current Opinion in Immunology* 16: 48–62.
- Jones JDG, Dangl JL. 2006. The plant immune system. *Nature* 444: 323–329.
- Jordan ND, Franklin FC, Franklin-Tong VE. 2000. Evidence for DNA fragmentation triggered in the self-incompatibility response in pollen of *Papaver rhoeas*. *Plant Journal* 23: 471–479.
- Kachroo A, Schopfer CR, Nasrallah ME, Nasrallah JB. 2001. Allele-specific receptor–ligand interactions in *Brassica* self-incompatibility. *Science* 293: 1824–1826.
- Kakita M, Shimamoto H, Murase K, Isogai A, Takayama S. 2007. Direct interaction between S-locus receptor kinase and M-locus protein kinase involved in *Brassica* self-incompatibility signalling. *Plant Biotechnology* 24: 185–190.
- Kaku H, Nishizawa Y, Ishii-Minami N, Akimoto-Tomiya C, Dohmae N, Takio N, Minami E, Shibuya N. 2006. Plant cells recognize chitin fragments for defense signaling through a plasma membrane receptor. *Proceedings of the National Academy of Sciences, USA* 103: 11086–11091.
- Kemp BP, Doughty J. 2007. S Cysteine-rich (SCR) binding domain analysis of the *Brassica* self-incompatibility S-locus receptor kinase. *New Phytologist* 175: 619–629.
- Liu J, Liu X, Dai L, Wang G. 2007. Recent progress in elucidating the structure, function and evolution of disease resistance genes in plants. *Journal of Genetics and Genomics* 34: 765–776.
- Matzinger P. 2002. The danger model: a renewed sense of self. *Science* 296: 301–305.
- Medzhitov R, Janeway CA. 2002. Decoding the patterns of self and nonself by the innate immune system. *Science* 296: 298–300.
- Meindl T, Boller T, Felix G. 2000. The bacterial elicitor flagellin activates its receptor in tomato cells according to the address-message concept. *Plant Cell* 12: 1783–1794.
- Mishima M, Takayama S, Sasaki K, Lee J, Kojima C, Isogai A, Shirakawa M. 2003. Structure of the male determinant factor for *Brassica* self-incompatibility. *Journal of Biological Chemistry* 278: 36389–36395.
- Mishina TE, Zeier J. 2007. Pathogen-associated molecular pattern recognition rather than development of tissue necrosis contributes to bacterial induction of systemic acquired resistance in Arabidopsis. *Plant Journal* 50: 500–513.
- Morillo SA, Tax FE. 2006. Functional analysis of receptor-like kinases in monocots and dicots. *Current Opinions in Plant Biology* 7: 391–399.
- Murase K, Shiba H, Iwano M, Che FS, Watanabe M, Isogai A, Takayama S. 2004. A membrane-anchored protein kinase involved in *Brassica* self-incompatibility signaling. *Science* 303: 1516–1519.
- Naithani S, Chookajorn T, Ripoll DR, Nasrallah JB. 2007. Structural modules for receptor dimerization in the S-locus receptor kinase extracellular domain. *Proceedings of the National Academy of Sciences, USA* 104: 12211–12216.
- Nasrallah JB. 2005. Recognition and rejection of self in plant self-incompatibility: comparisons to animal histocompatibility. *Trends in Immunology* 26: 412–418.
- Navarro L, Zipfel C, Rowland O, Keller I, Robatzek S, Boller T, Jones JDG. 2004. The transcriptional innate immune response to flg22; interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis. *Plant Physiology* 135: 113–128.
- Newbigin E, Vierstra RD. 2003. Sex and self-denial. *Nature* 423: 229–230.
- Nürnberger T, Brunner F, Kemmerling B, Piater L. 2004. Innate immunity in plants and animals: striking similarities and obvious differences. *Immunology Review* 198: 249–266.
- Nürnberger T, Kemmerling B. 2006. Receptor protein kinases – pattern recognition receptors in plant immunity. *Trends in Plant Science* 11: 519–522.

- Ohtake Y, Takahashi T, Komeda Y. 2000. Salicylic acid induces the expression of a number of receptor-like kinase genes in *Arabidopsis thaliana*. *Plant Cell Physiology* 41: 1038–1044.
- Pastuglia M, Roby D, Dumas C, Cock JM. 1997. Rapid induction by wounding and bacterial infection of an S gene family receptor-like kinase gene in *Brassica oleracea*. *Plant Cell* 9: 49–60.
- Pastuglia M, Swarup R, Rocher A, Saindrenan P, Roby D, Dumas C, Cock JM. 2002. Comparison of the expression patterns of two small gene families of S gene family receptor kinase genes during the defense response in *Brassica oleracea* and *Arabidopsis thaliana*. *Gene* 282: 215–225.
- Ron M, Avni A. 2004. The receptor for the fungal elicitor ethylene-inducing xylanase is a member of a resistance-like gene family in tomato. *Plant Cell* 16: 1604–1615.
- Sanabria NM, Dubery IA. 2006. Differential display profiling of the *Nicotiana* response to LPS reveals elements of plant basal resistance. *Biochemical and Biophysical Research Communications* 344: 1001–1007.
- Schopfer CR, Nasrallah ME, Nasrallah JB. 1999. The male determinant of self-incompatibility in *Brassica*. *Science* 286: 1697–1700.
- Shapiro JA. 2002. A 21st century view of evolution. *Journal of Biological Physics* 28: 745–764.
- Shimosato H, Yokota N, Shiba H, Iwano M, Entani T, Che FS, Watanabe M, Isogai A, Takayama S. 2007. Characterisation of the SP11/SCR high-affinity binding site involved in self/nonself recognition in *Brassica* self-incompatibility. *Plant Cell* 19: 107–117.
- Shiu S, Bleecker AB. 2001. Plant receptor-like kinase gene family: diversity, function and signalling. *Science Signal Transduction Knowledge Environment* 113: re22.
- Shiu S-H, Karlowski WM, Pan R, Tzeng Y-H, Mayer KFX, Li W-H. 2004. Comparative analysis of the receptor-like kinase family in Arabidopsis and rice. *Plant Cell* 16: 1220–1234.
- Silva NF, Stone SL, Christie LN, Sulaman W, Nazarian KA, Burnett LA, Arnoldo MA, Rothstein SJ, Goring DR. 2001. Expression of the S receptor kinase in self-compatible *Brassica napus* cv. Westar leads to the allele-specific rejection of self-incompatible *Brassica napus* pollen. *Molecular Genetics and Genomics* 265: 552–559.
- Stone SL, Anderson EM, Mullen RT, Goring DR. 2003. ARC1 is an E3 ubiquitin ligase and promotes the ubiquitination of proteins during the rejection of self-incompatible *Brassica* pollen. *Plant Cell* 15: 885–898.
- Stone SL, Arnoldo M, Goring DR. 1999. A breakdown of *Brassica* self-incompatibility in ARC1 antisense transgenic plants. *Science* 286: 1729–1731.
- Takahashi T, Mu JH, Gasch A, Chua NH. 1998. Identification by PCR of receptor-like protein kinases from Arabidopsis flowers. *Plant Molecular Biology* 37: 587–596.
- Takasaki T, Hatakeyama K, Suzuki G, Watanabe M, Isogai A, Hinata K. 2000. The S receptor kinase determines self-incompatibility in *Brassica* stigma. *Nature* 403: 913–916.
- Takayama S, Isogai A. 2005. Self-incompatibility in plants. *Annual Review of Plant Biology* 56: 467–489.
- Takayama S, Shiba H, Iwano M, Shimosato H, Che FS. 2000. The pollen determinant of self-incompatibility in *Brassica campestris*. *Proceedings of the National Academy of Sciences, USA* 97: 1920–1925.
- Takayama S, Shimosato H, Shiba H, Funato M, Che FE, Watanabe M, Iwano M, Isogaim A. 2001. Direct ligand–receptor complex interaction controls *Brassica* self-incompatibility. *Nature* 413: 534–538.
- Tameling WIL, Takken FLW. (in press). Resistance proteins: scouts of the plant innate immune system. *European Journal of Plant Pathology* (published online: 27 September 2007).
- Thomas SG, Franklin-Tong VE. 2004. Self-incompatibility triggers programmed cell death in *Papaver* pollen. *Nature* 429: 305–309.
- Thordal-Christensen H. 2003. Fresh insights into processes of nonhost resistance. *Current Opinions in Plant Biology* 6: 351–357.
- Tordai H, Banyani L, Patthy L. 1999. The PAN module: the N-terminal domains of plasminogen and hepatocyte growth factor are homologous with the apple domains of the prekallikrein family and with a novel domain found in numerous nematode proteins. *FEBS Letters* 461: 63–67.
- Torii KU. 2004. Leucine-rich repeat receptor kinases in plants: structure, function, and signal transduction pathways. *International Review of Cytology* 234: 1–46.
- Umehoto N, Kakitani M, Iwamatsu A, Yoshikawa M, Yamaoka N, Ishida I. 1997. The structure and function of a soybean beta-glucan-elicitor-binding protein. *Proceedings of the National Academy of Sciences, USA* 94: 1029–1034.
- Vanoosthuyse V, Tichtinsky G, Dumas C, Gaude T, Cock JM. 2003. Interaction of calmodulin, a sorting nexin and kinase-associated protein phosphatase, with the *Brassica oleracea* S locus receptor kinase. *Plant Physiology* 133: 919–929.
- Walker JC. 1993. Receptor-like protein kinase genes of *Arabidopsis thaliana*. *Plant Journal* 3: 451–456.
- Walker JC. 1994. Structure and function of the receptor-like kinases of higher plants. *Plant Molecular Biology* 26: 1599–1609.
- Wheeler MJ, Franklin-Tong VE. 2007. Specifying self-recognition: peptides lead the way. *New Phytologist* 175: 597–599.
- Yamaguchi Y, Pearce G, Ryan CA. 2006. The cell surface leucine-rich repeat receptor for AtPep1, an endogenous peptide elicitor in Arabidopsis, is functional in transgenic tobacco cells. *Proceedings of the National Academy of Sciences, USA* 103: 10104–10109.
- Yang CW, Gonzalez-Lamothe R, Ewan RA, Rowland O, Yoshioka H, Shenton M, Ye H, O'Donnell E, Jones JD, Sadanandom A. 2006. The E3 ubiquitin ligase activity of Arabidopsis plant U-Box17 and its functional tobacco homolog ACRE276 are required for cell death and defense. *Plant Cell* 18: 1084–1098.
- Zeng LR, Qu S, Bordeos A, Yang C, Baraoidan M, Yan H, Xie Q, Nahm BH, Leung H, Wang GL. 2004. Spotted leaf1, a negative regulator of plant cell death and defense, encodes a U-box/armadillo repeat protein endowed with E3 ubiquitin ligase activity. *Plant Cell* 16: 2795–2808.
- Zhang X. 1998. Leucine-rich repeat receptor-like kinases in plants. *Plant Molecular Biological Report* 16: 301–311.
- Zipfel C, Felix G. 2005. Plants and animals: a different taste for microbes? *Current Opinions in Plant Biology* 8: 353–360.
- Zipfel C, Kunze G, Chinchilla D, Caniard A, Jones JDG, Boller T, Felix G. 2006. Perception of the bacterial PAMP EF-Tu by the receptor ERF restricts *Agrobacterium*-mediated transformation. *Cell* 125: 749–760.
- Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JDG, Felix G, Boller T. 2004. Bacterial disease resistance in Arabidopsis through flagellin perception. *Nature* 428: 764–767.



About *New Phytologist*

- *New Phytologist* is owned by a non-profit-making **charitable trust** dedicated to the promotion of plant science, facilitating projects from symposia to open access for our Tansley reviews. Complete information is available at www.newphytologist.org.
- Regular papers, Letters, Research reviews, Rapid reports and both Modelling/Theory and Methods papers are encouraged. We are committed to rapid processing, from online submission through to publication 'as-ready' via *OnlineEarly* – our average submission to decision time is just 28 days. Online-only colour is **free**, and essential print colour costs will be met if necessary. We also provide 25 offprints as well as a PDF for each article.
- For online summaries and ToC alerts, go to the website and click on 'Journal online'. You can take out a **personal subscription** to the journal for a fraction of the institutional price. Rates start at £135 in Europe/\$251 in the USA & Canada for the online edition (click on 'Subscribe' at the website).
- If you have any questions, do get in touch with Central Office (newphytol@lancaster.ac.uk; tel +44 1524 594691) or, for a local contact in North America, the US Office (newphytol@ornl.gov; tel +1 865 576 5261).